

Routine Schedule

Age	Vaccine	Primary Doses	Booster Dose
6 months– 4 years	Pfizer– Infant/Toddler	1st Dose → 3-8 weeks [^] → 2nd Dose → ≥8 weeks → 3rd Dose	
6 months– 5 years	Moderna– Infant/Toddler	1st Dose → 4-8 weeks [^] → 2nd Dose	
5–11 years	Pfizer– Pediatric	1st Dose → 3-8 weeks [^] → 2nd Dose → ≥5 months →	Booster
6–11 years	Moderna– Pediatric	1st Dose → 4-8 weeks [^] → 2nd Dose	

Schedule if Moderately or Severely Immunocompromised

Age	Vaccine	Primary Doses	Booster Dose
6 months– 4 years	Pfizer– Infant/Toddler	1st Dose → 3 weeks → 2nd Dose → ≥8 weeks → 3rd Dose	
6 months– 5 years	Moderna– Infant/Toddler	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose	
5–11 years	Pfizer– Pediatric	1st Dose → 3 weeks → 2nd Dose → ≥4 weeks → 3rd Dose → ≥3 months →	Booster
6–11 years	Moderna– Pediatric	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose	

[^] An [8-week interval](#) may be preferable for some people, especially for males 12-39 years.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

COVID-19 Vaccine Timing–Adolescent/Adult

Routine Schedule

Age	Vaccine	Primary Doses	Booster Doses
12-17	Moderna– Adol/Adult		
12+	Pfizer/ Adol/Adult		
18+	Moderna– Adol/Adult		
18+	Janssen (J&J) Pfizer/Moderna preferred*		

^ An [8-week interval](#) may be preferable for some people, especially for males 12-39 years.

* Although use of mRNA COVID-19 vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

COVID-19 Vaccine Timing–Adolescent/Adult

Schedule if Moderately or Severely Immunocompromised

Age	Vaccine	Primary Doses	Booster Doses
12-17	Moderna– Adol/Adult		
12+	Pfizer/ Adol/Adult		<div> 1st Booster Ages 12-17: Pfizer 18+: Pfizer/ Moderna (mRNA preferred) or J&J* </div> <div> </div> <div> 2nd Booster Ages 12-17: Pfizer 18+: Moderna/ Pfizer </div>
18+	Moderna– Adol/Adult		
18+	Janssen (J&J) Pfizer/Moderna preferred*		

* Although use of mRNA COVID-19 vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.