WEEKLY MUMMY BERRY UPDATE

(2/14/2022 - 2/20/2022)



This weekly update provides information on the timing of apothecia (mushrooms) development from mummified overwintering berries in Washington's Skagit, Island, Snohomish, and Whatcom Counties.

The first table shows the percentages of floral and vegetative buds at different developmental stages and indicates when susceptible tissue is available on the plants. The second table shows the apothecial developmental stages from mummies and when ascospores are produced.

Development of mummies and host reflect conditions occurring in one field from each of the four counties that are currently being monitored. There may be differences in the stages of development in other blueberry fields in these same counties, in different counties, or for different blueberry cultivars.

Chakradhar Mattupalli

Berry and Potato Pathology Program WSU Mount Vernon Northwestern Washington Research and Extension Center Office phone: 360-848-6138

Email: c.mattupalli@wsu.edu

		Floral buds			Vegetative buds			
County	Cultivar						<1½"-7	
		Dormant (%)	Bud swell (%)	Bud break (%)	Dormant (%)	Bud swell (%)	Early green (%)	
Snohomish	Blueray	6.7	93.3	0	80	20	0	
	Bluecrop	66.7	33.3	0	100	0	0	
Skagit	Duke	0	100	0	100	0	0	
Island	Aurora	13.3	86.7	0	100	0	0	
	Legacy	0	90	10	0	90	10	
Whatcom	Rancocas	46.7	53.3	0	100	0	0	
	Bluecrop	10	90	0	70	23.3	6.7	
	Percentages represent data collected from 30 plants per each cultivar. Red indicates a susceptible stag infections.							

(Pictures: https://fieldguide.bcblueberry.com/crop-growth-stages/; Dr. Dalphy Harteveld; https://www.canr.msu.edu/blueberries/growing_blueberries/growth-stages)

County		94	S						
	Dormant (%)	Germination (%)	Emergence (%)	Differentiation (%)	Sporulation (%)	Finished (%)			
Snohomish	88	8	4	0	0	0			
Skagit	65	25	10	0	0	0			
Island	63	23	13	1	0	0			
Whatcom	66	22	11	1	0	0			
	Percentages represent data collected from 100 mummies in one field from each county. Red indicates release of ascospores.								