Indicators of Sacroiliac (SI) Joint Pain

It Hurts Right Here

Fortin Finger Sign

Simple, Reliable

Diagnostic Aid

Ask patient to point to location of primary pain:

– Below L5: consider SI joint

– Above L5: consider lumbar spine etiologies

Painful Activities

Sitting

– Weight on the unaffected side

Sit to stand

– Pain going from sitting to standing

– Midrise catch

– Use of hands/arms to assist with standing

Stepping up

– Pain going up stairs

– Pain with loading affected side

The iFuse Implant System® is intended for sacroiliac fusion for conditions including sacroiliac joint dysfunction that is a direct result of sacroiliac joint disruption and degenerative sacroiliitis. This includes conditions whose symptoms began during pregnancy or in the peripartum period and have persisted postpartum for more than 6 months. There are potential risks associated with the iFuse Implant System. It may not be appropriate for all patients and all patients may not benefit. For information about the risks, visit *www.si-bone.com/risks*

SI-BONE and *iFuse Implant System* are registered trademarks of SI-BONE, Inc. Santa Clara, CA. USA
©2018 SI-BONE, Inc. All rights reserved. Patents *www.si-bone.com* 9859.041918

SI-BONE®

Visit **si-bone.com** to learn more